


Chapter 3 Water and the Fitness of the Environment

Concept 3.1 The polarity of water molecules results in hydrogen bonding


- A water molecule is a polar molecule in which opposite ends of the molecule have opposite charges.
 - Because oxygen is more electronegative than hydrogen, the region around the oxygen atom has a partial negative charge.
 - The regions near the two hydrogen atoms have a partial positive charge.
 - The slightly negative regions of one water molecule are attracted to the slightly positive regions of nearby water molecules, forming hydrogen bonds.
 - Each water molecule can form hydrogen bonds with up to four neighbors.



Concept 3.2 Four emergent properties of water contribute to Earth's fitness for life

1. *Organisms depend on the cohesion of water molecules.*

- 1/20 as strong as covalent bonds.
- They form, break, and reform. Each hydrogen bond lasts only a few trillionths of a second.
- Each water can H bond to 4 other water at once
- Most water molecules are bonded to their neighbors, creating a high level of structure.
- Collectively, cause **cohesion**.
- Key role in the transport of water and dissolved nutrients against gravity in plants.
 - Roots, stems leaves are full of water-conducting vessels.
 - water molecules evaporate from a leaf,
 - Hydrogen bonds cause water molecules leaving to tug on molecules farther down.
 - upward pull is transmitted down to the roots.
 - **Adhesion**, clinging of one substance to another, contributes too, as water adheres to the wall of the vessels. (like capillary tube)



- **Surface tension**, a measure of the force necessary to stretch or break the surface of a liquid, is related to cohesion.
 - Surface water molecules resist stretching or breaking of the surface.
 - Water behaves as if covered by an invisible film.
 - Some animals can stand, walk, or run on water without breaking the surface.

2. Water moderates temperatures on Earth.

- Water stabilizes air temperatures by absorbing heat from warmer air and releasing heat to cooler air.
- Water can absorb or release relatively large amounts of heat with only a slight change in its own temperature.
- Atoms and molecules have **kinetic energy**, the energy of motion, because they are always moving.
 - The faster a molecule moves, the more kinetic energy it has.
- **Heat** is a measure of the *total* quantity of kinetic energy due to molecular motion in a body of matter.
- **Temperature vs. Heat**
 - Heat and temperature are related, but not identical.
 - Heat = a form of energy that can be transferred
 - Temp = a measure of the amount of heat. It is caused by the faster movement of molecules with more heat.
- In most biological settings, temperature is measured on the **Celsius scale** ($^{\circ}\text{C}$).
 - At sea level, water freezes at 0°C and boils at 100°C .
 - Human body temperature is typically 37°C ., room temp about 21°C
- A **calorie (cal)**.
 - One calorie is the amount of heat energy necessary to raise the temperature of one g of water by 1°C .
 - A calorie is released when 1 g of water cools by 1°C .
- In many biological processes, the **kilocalorie (kcal)** is more convenient.
 - A kilocalorie is the amount of heat energy necessary to raise the temperature of 1000 g of water by 1°C .
 - Food packaging lists energy in kcal (Calories with a "C")
- the **joule (J)** is a unit for tinier quantities of energy and is equivalent to 0.239 cal.
- Water stabilizes temperature because it has a high specific heat.
 - The **specific heat** of a substance is the amount of heat that must be absorbed or lost for 1 g of that substance to change its temperature by 1°C .
 - By definition, the specific heat of water is 1 cal per gram per degree Celsius or $1\text{ cal/g}^{\circ}\text{C}$.



Copyright 2012
Cengage Learning


- Water has a high specific heat compared to other substances.
 - Water resists changes in temperature because it absorbs or releases a relatively large quantity of heat for each degree of temperature change.
- Water's high specific heat is due to hydrogen bonding. Heat must be absorbed to break hydrogen bonds, and heat is released when hydrogen bonds form.
 - one calorie of heat causes relatively little change to the temperature of water - much of the energy is used to break hydrogen bonds, not speed up the movement of water molecules (temp).
- Water's high specific heat has effects that range from the level of the whole Earth to the level of individual organisms.
 - A large body of water can absorb a large amount of heat from the sun in daytime during the summer and yet warm only a few degrees.
 - At night and during the winter, the warm water will warm cooler air.
 - Therefore, ocean temperatures and coastal land areas have more stable temperatures than inland areas.
 - Living things are made primarily of water. Consequently, they resist changes in temperature better than they would if composed of a liquid with a lower specific heat.
- The transformation of a molecule from a liquid to a gas is called **vaporization** or **evaporation**.
 - This occurs when the molecule moves fast enough to overcome the attraction of other molecules in the liquid.
- **Heat of vaporization** is the quantity of heat that a liquid must absorb for 1 g of it to be converted from liquid to gas.
 - Water has a relatively high heat of vaporization, requiring about 580 cal of heat to evaporate 1 g of water at room temperature.
 - This is double the heat required to vaporize the same quantity of alcohol or ammonia.
 - This is because hydrogen bonds must be broken before a water molecule can evaporate from the liquid.
 - Water's high heat of vaporization moderates climate.
 - Much of the sun's heat absorbed by tropical oceans is used for evaporation of surface water.
 - As moist tropical air moves to the poles, water vapor condenses to form rain, releasing heat.
- As a liquid evaporates, the surface of the liquid that remains behind cools, a phenomenon called **evaporative cooling**.
 - This occurs because the most energetic molecules are the most likely to evaporate, leaving the lower-kinetic energy molecules behind.
- Evaporative cooling moderates temperature in lakes and ponds.
- Evaporation of sweat in mammals or evaporation of water from the leaves of plants prevents terrestrial organisms from overheating.

- Evaporation of water from the leaves of plants or the skin of humans removes excess heat.

Oceans and lakes don't freeze solid because ice floats.

- Water is unusual because it is less dense as a solid than as a cold liquid.

- Most materials contract as they solidify, but water expands.
- At temperatures above 4°C, water behaves like other liquids, expanding as it warms and contracting as it cools.
- Water begins to freeze when its molecules are no longer moving vigorously enough to break their hydrogen bonds.




- When water reaches 0°C, water becomes locked into a crystalline lattice, with each water molecule bonded to a maximum of four partners.
- As ice starts to melt, some of the hydrogen bonds break, and water molecules can slip closer together than they can while in the ice state.
- Ice is about 10% less dense than water at 4°C. Therefore, ice floats on the cool water below.
- This oddity has important consequences for life.
 - If ice sank, eventually all ponds, lakes, and even the ocean would freeze solid.
 - During the summer, only the upper few centimeters of the ocean would thaw.
 - Instead, the surface layer of ice insulates liquid water below, preventing it from freezing and allowing life to exist under the frozen surface.


Water is the solvent of life.

- A liquid that is a completely homogeneous mixture of two or more substances is called a **solution**.
 - A sugar cube in a glass of water will eventually dissolve to form a uniform solution of sugar and water.
 - The dissolving agent is the **solvent**, and the substance that is dissolved is the **solute**.
 - In our example, water is the solvent and sugar is the solute.
- In an **aqueous solution**, water is the solvent.
- Water is not a universal solvent, but it is very versatile because of the polarity of water molecules.
 - Water is an effective solvent because it readily forms hydrogen bonds with charged and polar covalent molecules.

- For example, when a crystal of salt (NaCl) is placed in water, each dissolved ion is surrounded by a sphere of water molecules, a **hydration shell**.



- Eventually, water dissolves all the ions, resulting in a solution with two solutes: sodium and chloride ions.
- Polar molecules are also soluble in water because they form hydrogen bonds with water.
- Even large molecules, like proteins, can dissolve in water if they have ionic and polar regions.
- Any substance that has an affinity for water is **hydrophilic** (*water-loving*).
 - These substances are dominated by ionic or polar bonds.
- Some hydrophilic substances do not dissolve because their molecules are too large.
 - For example, cotton is hydrophilic because cellulose, its major constituent, has numerous polar covalent bonds. However, its giant cellulose molecules are too large to dissolve in water.
 - Water molecules form hydrogen bonds with the cellulose fibers of cotton, allowing you to dry yourself with your cotton towel as the water is pulled into the towel.
- Substances that have no affinity for water are **hydrophobic** (*water-fearing*).
 - These substances are nonionic and have nonpolar covalent bonds.
 - Because there are no consistent regions with partial or full charges, water molecules cannot form hydrogen bonds with hydrophobic molecules.
 - Oils such as vegetable oil are hydrophobic because the dominant bonds, carbon-carbon and carbon-hydrogen, share electrons equally.
 - Hydrophobic molecules are major ingredients of cell membranes.
- Biological chemistry is “wet” chemistry with most reactions involving solutes dissolved in water.
- Chemical reactions depend on collisions of molecules and therefore on the concentrations of solutes in aqueous solution.
- We measure the number of molecules in units called **moles**.
- The actual number of molecules in a mole is called Avogadro’s number, 6.02×10^{23} .
- A mole is equal to the molecular weight of a substance in grams.
- The advantage of using moles as a measurement is that a mole of one substance has the same number of molecules as a mole of any other substance.
 - Measuring in moles allows scientists to combine substances in fixed ratios of molecules.
- In “wet” chemistry, we are typically combining solutions or measuring the quantities of materials in aqueous solutions.
 - The concentration of a material in solution is called its **molarity**.




- A one molar solution has one mole of a substance dissolved in one liter of solvent, typically water.
- To make a 1 molar (1M) solution of sucrose, we would slowly add water to 342 g of sucrose until the total volume was 1 liter and all the sugar was dissolved.

Concept 3.3 Dissociation of water molecules leads to acidic and basic conditions that affect living organisms

- Occasionally, a hydrogen atom participating in a hydrogen bond between two water molecules shifts from one molecule to the other.

- The hydrogen atom leaves its electron behind and is transferred as a single proton—a **hydrogen ion** (H^+).



- The water molecule that lost the proton is now a **hydroxide ion** (OH^-).

- The water molecule with the extra proton is now a **hydronium ion** (H_3O^+).

Hydronium ion (H_3O^+) Hydroxide ion (OH^-)

Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

- A simplified way to view this process is to say that a water molecule dissociates into a hydrogen ion and a hydroxide ion:
 - $H_2O \rightleftharpoons H^+ + OH^-$
- This reaction is reversible.
- Adding certain solutes, called acids and bases, disrupts the equilibrium and modifies the concentrations of hydrogen and hydroxide ions.
- The pH scale is used to describe how acidic or basic a solution is.

Organisms are sensitive to changes in pH.

- An **acid** is a substance that increases the hydrogen ion concentration in a solution.
 - When hydrochloric acid is added to water, hydrogen ions dissociate from chloride ions: $HCl \rightarrow H^+ + Cl^-$
 - Addition of an acid makes a solution more acidic.
- Any substance that reduces the hydrogen ion concentration in a solution is a **base**.
- Solutions with more OH^- than H^+ are basic solutions.
- Solutions with more H^+ than OH^- are acidic solutions.
- Solutions in which concentrations of OH^- and H^+ are equal are neutral solutions.
- Brackets ($[H^+]$ and $[OH^-]$) indicate the molar concentration of the enclosed substance.
- Adding a base does the opposite, increasing OH^- concentration and lowering H^+ concentration.
- The H^+ and OH^- concentrations of solutions can vary by a factor of 100 trillion or more.

- To express this variation more conveniently, the H^+ and OH^- concentrations are typically expressed via the pH scale.
 - The pH scale, ranging from 1 to 14, compresses the range of concentrations by employing logarithms.
 - $pH = -\log [H^+]$ or $[H^+] = 10^{-pH}$
 - In a neutral solution, $[H^+] = 10^{-7} M$, and the $pH = 7$.
- Values for pH *decline* as $[H^+]$ *increase*.
- While the pH scale is based on $[H^+]$, values for $[OH^-]$ can be easily calculated from the product relationship.
- The pH of a neutral solution is 7.
- Acidic solutions have pH values less than 7, and basic solutions have pH values greater than 7.
- Most biological fluids have pH values in the range of 6 to 8.
 - However, the human stomach has strongly acidic digestive juice with a pH of about 2.
- Each pH unit represents a tenfold difference in H^+ and OH^- concentrations.
 - A small change in pH actually indicates a substantial change in H^+ and OH^- concentrations.
- To maintain cellular pH values at a constant level, biological fluids have **buffers**.
- Buffers resist changes to the pH of a solution when H^+ or OH^- is added to the solution.
 - Buffers accept hydrogen ions from the solution when they are in excess and donate hydrogen ions when they have been depleted.
 - Buffers typically consist of a weak acid and its corresponding base.
 - One important buffer in human blood and other biological solutions is carbonic acid, which dissociates to yield a bicarbonate ion and a hydrogen ion.
 - The chemical equilibrium between carbonic acid and bicarbonate acts as a pH regulator. The equilibrium shifts left or right as other metabolic processes add or remove H^+ from the solution.

Acid precipitation threatens the fitness of the environment.

- **Acid precipitation** is a serious assault on water quality in some industrialized areas.
 - Uncontaminated rain has a slightly acidic pH of 5.6.
 - The acid is a product of the formation of carbonic acid from carbon dioxide and water.
- Acid precipitation occurs when rain, snow, or fog has a pH that is more acidic than 5.6.
- Acid precipitation is caused primarily by sulfur oxides and nitrogen oxides in the atmosphere.
 - These molecules react with water to form strong acids that fall to the surface with rain or snow.

- The major source of these oxides is the burning of fossil fuels (coal, oil, and gas) in factories and automobiles.
- The presence of tall smokestacks allows this pollution to spread from its site of origin to contaminate relatively pristine areas thousands of kilometers away.
 - In 2001, rain in the Adirondack Mountains of upstate New York had an average pH of 4.3.
- Thus, strong acidity can alter the structure of molecules and impact ecological communities.